

ASOTIN COUNTY 2007

32 CORE INDICATORS

Washington State has gathered data on these Local Public Health Indicators that were established in 2007. The results are shown on the tables below. Some data is not yet available.

1. Reported Chlamydia infections
2. Treated Chlamydia infections
3. Influenza vaccination (65 years of older)
4. Years of healthy life expected at age 20
5. Adult cigarette smoking
6. Adult physical activity
7. Adults overweight/obese
8. Adult fruit/vegetable consumption
9. Adult binge drinking
10. Adults with diabetes
11. Adult poor mental health
12. First trimester prenatal care
13. Maternal cigarette smoking
14. Teen birth rate
15. Low birth rate
16. Teen physical activity
17. Teen cigarette smoking
18. Teens overweight
19. Teen alcohol use
20. Childhood unintentional injury hospitalizations
21. Adults with unmet medical need
22. Adults with personal healthcare provider
23. Adult dental care
24. Adult preventive cancer screening – Breast
25. Adult preventive cancer screening – Cervical
26. Adult preventive cancer screening – Colorectal
27. Adults with health insurance
28. Children with health insurance
29. Children's immunization status
30. Permitted solid waste facilities
31. Food inspections with > 35 Critical Violations
32. On-Site System failures with Corrective Action within two weeks

Asotin County Health District

<i>Category:</i> Communicable Disease			Jurisdiction		State		US
Indicator Name	Year	Unit of Measure	Estimate	95% Confidence Interval	Estimate	95% Confidence Interval	Estimate
Reported Chlamydia infections ¹	2004 - '06	rate per 100,000	1817	(3276, 2270)	2234	(2208, 2260)	2805**
Treated Chlamydia infections ¹	2004 - '06	%	91	(84, 97)	93	(92, 94)	N/A*
Influenza vaccination (65 years or older) ²	2004 - '06	%	73	(67, 79)	69	(68, 70)	69
Children's immunization status	N/A*	%	N/A*	N/A*	N/A*	N/A*	N/A*

*N/A - data not available

**US year: 2003 - '05

Missing Data

†10%-19% missing data
 ††20% or more missing data

Suppressed Data

‡Data suppressed due to small numbers (less than 5 records per year or relative standard error >30%)


Source

¹State and local data: DOH, Sexually Transmitted Disease Registry
 National data: CDC, Division of Sexually Transmitted Disease Prevention
²State and local data: DOH, Behavioral Risk Factor Surveillance System
 National data: CDC, Behavioral Risk Factor Surveillance System

1. Rate of reported Chlamydia infections (women 15 – 24 years):

Indicator: **Reported Chlamydia infections**

2004 - 2006



Indicator: Rate of reported Chlamydia infections per 100,000 women ages 15 to 24

Key

- better than state*
- similar to state
- state
- worse than state*

*Statistically significant difference (p < .05)

Missing Data

- † 10%-19% missing data
- †† 20% or more missing data

Suppressed Data

Data for the following health jurisdictions have been suppressed due to small numbers (less than 5 records per year or relative standard error > 30%):

- Columbia
- Wahkiakum
- Garfield

Technical Notes


Sources

State and local data: DOH, Sexually Transmitted Disease Registry

2. % of reported adequate Chlamydia treatment for females 15 – 24 years:

Indicator: **Treated Chlamydia infections**

2004 - 2006



Indicator: Percent of reported Chlamydia infections that received treatment in women ages 15 to 24

Key

- better than state*
- similar to state
- state
- worse than state*

*Statistically significant difference (p< .05)

Missing Data

† 10%-19% missing data
 †† 20% or more missing data

Suppressed Data

Data for the following health jurisdictions have been suppressed due to small numbers (less than 5 records per year or relative standard error > 30%):
 - Columbia - Wahkiakum
 - Garfield


Technical Notes

Sources

State and local data: DOH, Sexually Transmitted Disease Registry

3. Influenza vaccine during previous year for 65+ years:

Indicator: Influenza vaccination (65 years or older)



Indicator: Percent of adults age 65 or older who report receiving a flu shot during the past 12 months

Key

- better than state*
- similar to state
- state
- worse than state*

*Statistically significant difference ($p < .05$)

Missing Data

† 10%-19% missing data

†† 20% or more missing data

Suppressed Data

Data for the following health jurisdictions have been suppressed due to small numbers (less than 5 records per year or relative standard error > 30%):
- none

Technical Notes

Sources

State and local data: DOH, Behavioral Risk Factor Surveillance System

<i>Category:</i> Prevention and Health Promotion			Jurisdiction		State		US
Indicator Name	Year	Unit of Measure	Estimate	95% Confidence Interval	Estimate	95% Confidence Interval	Estimate
Years of healthy life expected at age 20 ¹	2003 - '05	number of years	49	(46, 51)	51	(50, 51)	48**
Adult cigarette smoking ²	2004 - '06	%	24	(21, 29)	18	(17, 18)	20
Adult physical activity ²	2003 & '05	%	60	(55, 66)	63	(63, 64)	58
Adults overweight/obese ²	2004 - '06	%	58	(53, 62)	59	(59, 60)	61
Adult fruit/vegetable consumption ²	2003 & '05	%	24	(20, 29)	24	(24, 25)	24
Adult binge drinking ²	2006	%	22	(17, 28)	14	(13, 15)	15
Adults with diabetes ²	2004 - '06	%	8	(6, 10)	7	(6, 7)	8
Adult poor mental health ²	2004 - '06	%	10	(8, 13)	10	(10, 10)	10

*N/A - data not available

**US year: 2003

Source

¹State and local data: DOH, Behavioral Risk Factor Surveillance System and Death Certificates
National data: CDC, Behavioral Risk Factor Surveillance System combined with National Life Expectancy data

²State and local data: DOH, Behavioral Risk Factor Surveillance System
National data: CDC, Behavioral Risk Factor Surveillance System

Missing Data

†10%-19% missing data
††20% or more missing data

Suppressed Data


‡Data suppressed due to small numbers (less than 5 records per year or relative standard error >30%)

Technical Notes

4. Expected years of healthy life at age 20.

Indicator: Years of healthy life expected at age 20

2003 - 2005



Indicator: Additional years a 20 year-old is expected to live in good, very good, or excellent health

Key

- better than state*
- similar to state
- state
- worse than state*

*Statistically significant difference (p < .05)

Missing Data

†10%-19% missing data

†† 20% or more missing data

Suppressed Data

Data for the following health jurisdictions have been suppressed due to small numbers (less than 5 records per year or relative standard error > 30%):
- none


Technical Notes

Sources

State and local data: DOH, Behavioral Risk Factor Surveillance System and Death Certificates

5. % of adults who report smoking everyday or some days.

Indicator: **Adult cigarette smoking**



Indicator: Percent of adults age 18 or older who smoked at least 100 cigarettes in their lifetime and are current smokers

Key

- better than state*
- similar to state
- state
- worse than state*

*Statistically significant difference ($p < .05$)

Missing Data

- † 10%-19% missing data
- †† 20% or more missing data

Suppressed Data

Data for the following health jurisdictions have been suppressed due to small numbers (less than 5 records per year or relative standard error > 30%):
- none

Technical Notes


Sources

State and local data: DOH, Behavioral Risk Factor Surveillance System

6. % of adults who report meeting moderate or vigorous physical activity.

Indicator: **Adult physical activity**

2003 & 2005



Indicator: Percent of adults age 18 or older who report moderate physical activity (30 minutes a day 5 times a week) or vigorous activity (20 minutes a day 3 times a week) in work or leisure

Key

- better than state*
- similar to state
- state
- worse than state*

*Statistically significant difference (p < .05)

Missing Data

†10%-19% missing data
 †† 20% or more missing data

Suppressed Data

Data for the following health jurisdictions have been suppressed due to small numbers (less than 5 records per year or relative standard error > 30%):
 - none

Technical Notes

Percent


Sources

State and local data: DOH, Behavioral Risk Factor Surveillance System

7. % of adults who are obese and overweight – BMI.

Indicator: **Adult overweight/obese**

2004 - 2006



Indicator: Percent of adults age 18 or older who have body mass index greater than 25

Key

- better than state*
- similar to state
- state
- worse than state*

*Statistically significant difference (p < .05)

Missing Data

†10%-19% missing data
 †† 20% or more missing data

Suppressed Data

Data for the following health jurisdictions have been suppressed due to small numbers (less than 5 records per year or relative standard error > 30%):
 - none

Technical Notes

Percent


Sources

State and local data: DOH, Behavioral Risk Factor Surveillance System

8. % of adults who report eating fruits and vegetables 5 or more times per day.

Indicator: **Adult fruit/vegetable consumption**

2003 & 2005



Indicator: Percent of adults age 18 or older who report eating fruits and vegetables 5 or more times per day

Key

- better than state*
- similar to state
- state
- worse than state*

*Statistically significant difference ($p < .05$)

Missing Data

- † 10%-19% missing data
- †† 20% or more missing data

Suppressed Data

Data for the following health jurisdictions have been suppressed due to small numbers (less than 5 records per year or relative standard error > 30%):
- none


Technical Notes

Sources

State and local data: DOH, Behavioral Risk Factor Surveillance System

9. % of adults who report binge drinking on one or more occasion in the past 30 days.

Indicator: **Adult binge drinking**



Indicator: Percent of adults age 18 or older who report binge drinking (5 drinks for men; 4 drinks for women) on at least 1 occasion in last 30 days

Key

- better than state*
- similar to state
- state
- worse than state*

*Statistically significant difference ($p < .05$)

Missing Data

† 10%-19% missing data
 †† 20% or more missing data

Suppressed Data

Data for the following health jurisdictions have been suppressed due to small numbers (less than 5 records per year or relative standard error > 30%):
 - none

Technical Notes


Sources

State and local data: DOH, Behavioral Risk Factor Surveillance System

10. % of adults who report diagnosis of diabetes.

Indicator: **Adults with diabetes**

2004 - 2006



Indicator: Percent of adults age 18 or older who have ever been told by a doctor that they have diabetes

Key

- better than state*
- similar to state
- state
- worse than state*

*Statistically significant difference ($p < .05$)

Missing Data

† 10%-19% missing data
 †† 20% or more missing data

Suppressed Data

Data for the following health jurisdictions have been suppressed due to small numbers (less than 5 records per year or relative standard error > 30%):
 - none

Technical Notes

Percent


Sources

State and local data: DOH, Behavioral Risk Factor Surveillance System

11. % of adults who report 14 or more days of poor mental health in the past month.

Indicator: **Adult poor mental health**

2004 - 2006



Indicator: Percent of adults age 18 or older who report 14 or more days of poor mental health in the past month

Key

- better than state*
- similar to state
- state
- worse than state*

* Statistically significant difference (p < .05)

Missing Data

† 10%-19% missing data
 †† 20% or more missing data

Suppressed Data

Data for the following health jurisdictions have been suppressed due to small numbers (less than 5 records per year or relative standard error > 30%):
 - none

Technical Notes

Sources

State and local data: DOH, Behavioral Risk Factor Surveillance System

--	--	--

<i>Category: Maternal and Child Health</i>			Jurisdiction				
Indicator Name	Year	Unit of Measure	Estimate	95% Confidence Interval	Estimate	95% Confidence Interval	Estimate
First trimester prenatal care ¹	2003 - '05	%	78	(72, 85)	80†	(80, 80)	84**
Maternal cigarette smoking ¹	2003 - '05	%	23	(20, 27)	10	(10, 11)	10**
Teen birth rate ¹	2003 - '05	rate per 1,000	21	(15, 30)	15	(15, 16)	22**
Low birth weight ¹	2003 - '05	%	6	(5, 9)	5	(5, 5)	6**
Teen physical activity ²	2006	%	50	(39, 60)	42	(40, 44)	36***
Teen cigarette smoking ²	2006	%	20	(15, 27)	15	(14, 16)	21***
Teens overweight ²	2006	%	24	(16, 34)	25	(22, 27)	29***
Teen alcohol use ²	2006	%	43	(36, 50)	33	(31, 34)	42***
Childhood unintentional Injury hospitalizations	2002 - '04	rate per 100,000	N/A‡	N/A‡	199	(194, 203)	N/A*

*N/A - data not available
 **US year: 2004
 ***US year: 2005



Missing Data

10%-19% missing data
 †20% or more missing data

Suppressed Data

‡Data suppressed due to small numbers (less than 5 records per year or relative standard error >30%)
 ††Data suppressed due to small numbers (less than 5 records per year or less than 15 surveys per school), survey response rate <40%, or relative standard error >30%


Source

¹State and local data: DOH, Birth Certificates
 National data: CDC, National Vital Statistics System
²State and local data: DOH, Healthy Youth Survey
 National data: CDC, Youth Risk Behavior Surveillance System
³State and local data: DOH, Comprehensive Hospital Abstract Reporting System, including all acute care hospitals in Washington and Oregon, Madigan Army Hospital, Bremerton Naval Hospital and all VA Hospitals in the U.S.
 National data: Comparable national data are not available

12. % of women who received prenatal care during 1st trimester.

Indicator: **First trimester prenatal care**

2003 - 2005



Indicator: Percent of women giving birth who received prenatal care starting in the first trimester of pregnancy

Key

- better than state*
- similar to state
- state
- worse than state*

*Statistically significant difference (p < .05)

Missing Data

- † 10%-19% missing data
- †† 20% or more missing data

Suppressed Data

Data for the following health jurisdictions have been suppressed due to small numbers (less than 5 records per year or relative standard error > 30%):
- none

Technical Notes


Sources

State and local data: DOH, Birth Certificates

13. % of pregnant women who smoke during 2nd/3rd trimester of pregnancy.

Indicator: **Maternal cigarette smoking**

2003 - 2005



Indicator: Percent of women giving birth who smoked any time during pregnancy

Key

- better than state*
- similar to state
- state
- worse than state*

*Statistically significant difference (p < .05)

Missing Data

- † 10%-19% missing data
- †† 20% or more missing data

Suppressed Data

Data for the following health jurisdictions have been suppressed due to small numbers (less than 5 records per year or relative standard error > 30%):
- Garfield

Technical Notes

Percent


Sources

State and local data: DOH, Birth Certificates

14. Birth rate for females (age 15 – 17).

Indicator: **Teen birth rate**

2003 - 2005



Indicator: Rate of live born infants per 1,000 women ages 15 to 17

Key

- better than state*
- similar to state
- state
- worse than state*

*Statistically significant difference ($p < .05$)

Missing Data

- † 10%-19% missing data
- †† 20% or more missing data

Suppressed Data

Data for the following health jurisdictions have been suppressed due to small numbers (less than 5 records per year or relative standard error > 30%):

- Columbia
- Lincoln - Skamania
- Garfield
- San - Wahkiakum
- Juan

Technical Notes


Sources

State and local data: DOH, Birth Certificates

15. % low birth weight rate among singletons (< 2500g, 3 year average).

Indicator: **Low birth weight**

2003 - 2005



Indicator: Percent of live born singleton births with birth weight less than 2,500 grams

Key

- better than state*
- similar to state
- state
- worse than state*

*Statistically significant difference (p < .05)

Missing Data

- † 10%-19% missing data
- †† 20% or more missing data

Suppressed Data

Data for the following health jurisdictions have been suppressed due to small numbers (less than 5 records per year or relative standard error > 30%):

- Columbia
- Wahkiakum
- Garfield

Technical Notes


Percent

Sources

State and local data: DOH, Birth Certificates

16. % of 10th graders who report having not met recommendations for vigorous physical activity.

Indicator: **Teen physical activity**



Indicator: Percent of 10th graders who report physical activity 60 minutes a day, 5 or more days a week

Key

- better than state*
- similar to state
- state
- worse than state*

*Statistically significant difference ($p < .05$)

Missing Data

- † 10%-19% missing data
- †† 20% or more missing data

Suppressed Data

Data suppressed due to small numbers (less than 5 records per year or less than 15 surveys per school), survey response rate <40%, or relative standard error >30%:

- Clallam
- Northeast Tri County
- Grant


Technical Notes

Sources

State and local data: DOH, Healthy Youth Survey

17. Teen cigarette smoking

Indicator: Teen cigarette smoking



Indicator: Percent of 10th graders who report smoking cigarettes in the last 30 days

Key

- better than state*
- similar to state
- state
- worse than state*

*Statistically significant difference ($p < .05$)

Missing Data

- †10%-19% missing data
- †† 20% or more missing data

Suppressed Data

Data suppressed due to small numbers (less than 5 records per year or less than 15 surveys per school), survey response rate <40%, or relative standard error >30%:

- Clallam
- Columbia
- Garfield - Northeast Tri County
- Grant
- Wahkiakum


Technical Notes

Sources

State and local data: DOH, Healthy Youth Survey

18. 8% of 10th graders who are overweight – BMI.

Indicator: **Teens overweight**



Indicator: Percent of 10th graders in the top 5% body mass index by reported height and weight, based on CDC growth charts

Key

- better than state*
- similar to state
- state
- worse than state*

*Statistically significant difference ($p < .05$)

Missing Data

- †10%-19% missing data
- †† 20% or more missing data

Suppressed Data

Data suppressed due to small numbers (less than 5 records per year or less than 15 surveys per school), survey response rate <40%, or relative standard error >30%:

- Clallam
- Columbia
- Northeast Tri County
- Garfield - Wahkiakum
- Grant


Technical Notes

Sources

State and local data: DOH, Healthy Youth Survey

19. % of 10th graders who report alcohol consumption in the past 30 days.

Indicator: Teen alcohol use



Indicator: Percent of 10th graders who report alcohol consumption in the past 30 days

Key

- better than state*
- similar to state
- state
- worse than state*

*Statistically significant difference ($p < .05$)

Missing Data

† 10%-19% missing data
 †† 20% or more missing data

Suppressed Data

Data suppressed due to small numbers (less than 5 records per year or less than 15 surveys per school), survey response rate <40%, or relative standard error >30%:

- Clallam
- Grant
- Garfield
- Northeast Tri County

Technical Notes


Sources

State and local data: DOH, Healthy Youth Survey

20. Unintentional injury hospitalizations (age 0 – 17, 3 year average).

Indicator: **Childhood unintentional injury hospitalizations**

2002 - 2004



Indicator: Rate of hospitalization for unintentional injury per 100,000 children ages 0 to 17

Key

- better than state*
- similar to state
- state
- worse than state*

*Statistically significant difference ($p < .05$)

Missing Data

† 10%-19% missing data
 †† 20% or more missing data

Suppressed Data

Data for the following health jurisdictions have been suppressed due to small numbers (less than 5 records per year or relative standard error > 30%):

- Asotin
- Columbia
- Garfield - Skamania
- San Juan
- Wahkiakum

Technical Notes

Sources

State and local data: DOH, Comprehensive Hospital Abstract Reporting System, including all acute care hospitals in Washington and Oregon, Madigan Army Hospital, Bremerton Naval Hospital and all VA Hospitals in the U.S.

<i>Category: Access to Care</i>			Jurisdiction		State		US
Indicator Name	Year	Unit of Measure	Estimate	95% Confidence Interval	Estimate	95% Confidence Interval	Estimate
Adults with unmet medical need	2004 - '06	%	18	(15, 21)	13	(13, 14)	14
Adults with personal healthcare provider	2004 - '06	%	79	(75, 83)	78	(78, 79)	80
Adult dental care	2004 & '06	%	59	(54, 64)	71	(71, 72)	70
Adult preventive cancer screening - Breast	2004 & '06	%	71	(64, 77)	79	(79, 80)	80
Adult preventive cancer screening-Cervical	2004 & '06	%	77	(72, 82)	79	(78, 80)	81
Adult preventive cancer screening-Colorectal	2004 & '06	%	64	(58, 70)	62	(61, 63)	N/A*
Adults with health insurance	2004 - '06	%	78	(74, 82)	83	(82, 83)	81
Children with health insurance	N/A*	%	N/A*	N/A*	N/A*	N/A*	N/A*

*N/A - data not available

Missing Data

†10%-19% missing data
 ††20% or more missing data

Suppressed Data


‡Data suppressed due to small numbers (less than 5 records per year or relative standard error >30%)

Source

State and local data: DOH, Behavioral Risk Factor Surveillance System
 National data: CDC, Behavioral Risk Factor Surveillance System

21. % of adults in households who report unmet medical need due to cost.

Indicator: **Adults with unmet medical need**



Indicator: Percent of adults age 18 or older who report needing to see a doctor within the past year but could not due to cost

Key

- better than state*
- similar to state
- state
- worse than state*

* Statistically significant difference ($p < .05$)

Missing Data

- † 10%-19% missing data
- †† 20% or more missing data

Suppressed Data

Data for the following health jurisdictions have been suppressed due to small numbers (less than 5 records per year or relative standard error > 30%):
- none

Technical Notes


Percent

Sources

State and local data: DOH, Behavioral Risk Factor Surveillance System

22. % of adults who report having a Personal Doctor or Health Care Provider.

Indicator: **Adults with personal healthcare provider**



Indicator: Percent of adults age 18 or older who report having a personal doctor or healthcare provider

Key

- better than state*
- similar to state
- state
- worse than state*

*Statistically significant difference ($p < .05$)

Missing Data

- † 10%-19% missing data
- †† 20% or more missing data

Suppressed Data

Data for the following health jurisdictions have been suppressed due to small numbers (less than 5 records per year or relative standard error > 30%):
- none

Technical Notes


Percent

Sources

23. % of adults who report having visited dentist in past year.

Indicator: **Adult dental care**

2004 & 2006



Indicator: Percent of adults age 18 or older who report visiting a dentist, dental hygienist or dental clinic within the past year

Key

- better than state*
- similar to state
- state
- worse than state*

*Statistically significant difference (p < .05)

Missing Data

- † 10%-19% missing data
- †† 20% or more missing data

Suppressed Data

Data for the following health jurisdictions have been suppressed due to small numbers (less than 5 records per year or relative standard error > 30%):
- none


Technical Notes

Sources

24. % of adults who report receiving preventative cancer screenings – breast, cervical, colorectal.

Indicator: **Adult preventive cancer screening - Breast**

2004 & 2006



Indicator: Percent of women age 50 or older who report receiving a mammogram within the past 2 years

Key

- better than state*
- similar to state
- state
- worse than state*

* Statistically significant difference ($p < .05$)

Missing Data

† 10%-19% missing data
 †† 20% or more missing data

Suppressed Data

Data for the following health jurisdictions have been suppressed due to small numbers (less than 5 records per year or relative standard error > 30%):
 - none

Technical Notes

Percent


Sources

State and local data: DOH, Behavioral Risk Factor Surveillance System

25.

Indicator: **Adult preventive cancer screening - Cervical**

2004 & 2006



Indicator: Percent of women age 18 or older who report receiving a Pap smear test within the past 3 years

Key

- better than state*
- similar to state
- state
- worse than state*

*Statistically significant difference (p < .05)

Missing Data

- † 10%-19% missing data
- †† 20% or more missing data

Suppressed Data

Data for the following health jurisdictions have been suppressed due to small numbers (less than 5 records per year or relative standard error > 30%):
- none

Technical Notes


Sources

Percent

26.

Indicator: **Adult preventive cancer screening - Colorectal**

2004 & 2006



Indicator: Percent of adults age 50 or older who had a blood stool test in the past year, sigmoidoscopy in the past 5 years, or colonoscopy in the past 10 years

Key

- better than state*
- similar to state
- state
- worse than state*

*Statistically significant difference (p < .05)

Missing Data

- † 10%-19% missing data
- †† 20% or more missing data

Suppressed Data

Data for the following health jurisdictions have been suppressed due to small numbers (less than 5 records per year or relative standard error > 30%):
- none


Technical Notes

Sources

27. % of adults who report having health insurance.

Indicator: **Adults with health insurance**

2004 - 2006



Indicator: Percent of adults age 18-64 who currently report having health care insurance

Key

- better than state*
- similar to state
- state
- worse than state*

*Statistically significant difference ($p < .05$)

Missing Data

- † 10%-19% missing data
- †† 20% or more missing data

Suppressed Data

Data for the following health jurisdictions have been suppressed due to small numbers (less than 5 records per year or relative standard error > 30%):
- none

Technical Notes

Sources

State and local data: DOH, Behavioral Risk Factor Surveillance System

No data available for the following indicators:

28. % of children who are reported as having insurance.
29. % of Children who are Adequately Immunized at 24 months of age
30. % of Permitted Solid Waste Facilities in Compliance with Permit Conditions
31. % of Inspections of Permanent Food Establishments with 35 or > Critical Violations
32. % of Identified on-Site Sewage System Failures Initiated with Corrected Action within 2 weeks

